

# Agra Base

Reg. No. B6057 Act. No. 36 of 1947

A POWDER PLANT NUTRIENT / 'N POEIER PLANTVOEDINGSTOF

A fertilizer containing micro-elements.

'n Voedingsproduk wat mikro-elemente bevat.

## FERTILIZER GROUP 2 / MISSTOF GROEP 2

### PRODUCT COMPOSITION / PRODUK SAMESTELLING

Zn: 61 g/kg  
Cu: 17.4 g/kg  
Mn: 44 g/kg  
B: 18 g/kg  
Mo: 0.9 g/kg

NET MASS

20 kg

NETTO MASSA

### REGISTRATION HOLDER / REGISTRASIEHOUER

AgraForUm South Africa (Pty) Ltd

Reg. no: 2004/022326/07

P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330  
5 Ednau Small Holdings, Abrahamskraal road, Mooiwater, Bloemfontein, 9338

### DISTRIBUTOR / VERSPREIDER

AgraForUm South Africa (Pty) Ltd

Reg. no: 2004/022326/07

P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330

Batch number / Date of manufacturing:

Lot nommer / Datum van vervaardiging:

### PRECAUTION

- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.

### GENERAL

- **Agra Base** is a fertilizer intended as a supplement to normal fertilizer programs.
- Foliar spraying should preferably be done during the cooler periods of the day (ideal 20°C; < 40°C), as well as when the relative humidity is on its highest (before 9 am in the morning and after 4 pm in the afternoon).
- Wind speed should be < 8 km/h.
- Do not apply to plants when plants are wilted or under stress.

### APPLICATION RATES

CROPS	APPLICATION RATE	RECOMMENDATIONS
Row Crops	Foliar spray 0,2% concentration (maximum 4 kg/ha) Via irrigation system or soil 1-15 kg/ha	Use minimum 200 L water/ha for foliar application. Optimum results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Vegetable crops	Foliar spray 0,2% concentration (maximum 4 kg/ha) Via irrigation system or soil 1-15 kg/ha	Use minimum 300 L water/ha for foliar application. Optimum results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Fruit Tree crops	Foliar spray 0,2% concentration (maximum 4 kg/ha) Via irrigation system or soil 1-15 kg/ha	Once monthly during season as required based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.

### WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time, and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

AgraForum

Agra Base



VR PRINT (031) 569 1235  
ATTACH-A-LEAFLET®