



Reg. No. B5762 Act. No. 36 of 1947

# Agra Co

# 10L

A POWDER PLANT NUTRIENT  
A fertilizer containing Cobalt.

'N POEIER PLANTVOEDINGSTOF  
'n Voedingsprodukt wat Kobalt bevat.

FERTILIZER GROUP

# 2

MISSTOF GROEP

## ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Cobalt \_\_\_\_\_ 200 g/kg \_\_\_\_\_ Kobalt



REGISTRATION HOLDER / REGISTRASIE HOUER

AgraForum South Africa (Pty) Ltd | Reg. no: 2004/022326/07

P.O. Box 38910, Langenhovenpark, Bloemfontein, 9330 | [www.agraforum-sa.com](http://www.agraforum-sa.com)

Batch Number | Date of manufacturing  
Lot Nommer | Datum van vervaardiging



## GENERAL

- First dissolve the product in clean container with water before addition to spray tank.
- Foliar spraying should preferably be done during the cooler periods of the day (ideal 20 °C; < 40°C), as well as when the relative humidity is on its highest.
- Wind speed should be < 8 km/h.
- Do not apply to plants when plants are wilted or under stress.
- Please use a buffering agent to buffer pH between 5 and 6.
- Please use a suitable water-soluble wetting agent and penetrant to improve wetting, sticking and penetrating properties.
- Try to use highest water volume possible per hectare.

## COMPATIBILITY

- Always verify compatibility when mixed with other agricultural remedies (jar test as well as first spraying a small area to check for phytotoxicity).
- Always test the product first on a small area to check for phytotoxicity especially if higher rates are used.

## APPLICATION RATES

Crop	Dosage rate	Remark
Grain-, cereal-, field and forage legume crops	Foliar spray 100-150 g/ha Via irrigation system 100-200 g/ha In-furrow application 100-150 g/ha	Use minimum 200 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-5 leaf stage based on soil and/or leaf analysis. For forage legumes apply at least 2 weeks before cutting. Do not exceed nutrient requirement of the crop.
Vegetable crops	Foliar spray 50-100 g/ha Via irrigation system 100-200 g/ha	Use minimum 300 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-5 leaf stage and repeat application as required (at least 2-3 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Fruit trees & Vines	Foliar spray 50-100 g/ha Via irrigation system 100-200 g/ha	Two to three applications per season 2-3 weeks after full bloom with last application 21 days before harvest. Apply as required based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.

## WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

## PRECAUTION

- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.
- Product contains Cobalt (Co).