



Reg. No. M60 Act. No. 36 of 1947

ComCat[®] MAX WP

500g

A BIOSTIMULANT
Enhancing plant growth and yield.

'N BIOSTIMULANT
Verbeter plant groei en opbrengs.

FERTILIZER GROUP

3

MISSTOF GROEP

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

24-Epibrassinolide _____ 10.5 mg/kg _____ 24-Epibrassinolied



Agraforum

REGISTRATION HOLDER / REGISTRASIE HOUER

AgraForUm South Africa (Pty) Ltd | Reg. no: 2004/022326/07
5 Ednau Small Holdings, Abramskraal road, Mooiwater, Bloemfontein 9338
Tel: 051 011 3997

DISTRIBUTOR / VERSPREIDER

AgraForUm South Africa (Pty) Ltd | Reg. no: 2004/022326/07
5 Ednau Small Holdings, Abramskraal road, Mooiwater, Bloemfontein 9338

Batch Number | Date of manufacturing
Lot Nommer | Datum van vervaardiging



PRECAUTIONS

- Its misuse may cause damage to crops.
- Mild irritant to eyes. Wear protective clothing. In case of eye contact, wash with clean running water.
- Do not eat, drink or smoke whilst mixing or applying.
- Store in a cool (below 24°C) and dry place.

GENERAL

- ComCat® MAX WP is water-insoluble and non-toxic to plants when applied as directed.
- For best results apply ComCat® according to recommendations.
- Applied via the irrigation system.
- Before using ComCat® MAX WP, consult an agriculturalist or the manufacturer on the use and compatibility with other products.

DIRECTIONS FOR USE

- Premix the prescribed amount of ComCat® MAX WP in 5-10 L of water. Add the required amount of Aqua-Wet® at 10ml/100L water (water volume of injection tank) to the ComCat® MAX WP mixture and stir thoroughly. Add AnnGro® at required amount (240 ml/ha) and let mixture stand for 30 minutes before addition to injection tank of irrigation system. Add ZumSil® at recommended rate to injection tank. Ensure thorough mixing during application. For best results, do not let prepared mixture stand overnight. The use of the correct adjuvant is recommended for better results.

METHOD OF APPLICATION

Crop	Time of Application	ComCat® MAX WP dosage (g/ha)	ZumSil® dosage (L/ha)
Maize	9-15 leaves unfolded	50	1
Wheat	Growth stage 8 or 16 (Joubert)	100	1

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

VOORSORGMATREËLS

- Wangebruik kan gewasse beskadig.
- Mag irriterend wees vir die oë. Dra beskermende klere. In die geval van oogkontak, was met skoon lopende water.
- Moet nie eet, drink of rook gedurende meng of aanwending nie.
- Berg in 'n koel (laer as 24°C) en droë area.

ALGEMEEN

- ComCat® MAX WP is nie oplosbaar in water nie en nie-toksies vir plante wanneer toegedien soos voorgeskryf.
- Vir beste resultate dien ComCat® MAX WP toe volgens aanbeveling.
- Toegedien d.m.v. besproeiingsstelsel.
- Voordat ComCat® MAX WP gebruik word, raadpleeg 'n landboukundige of die vervaardiger rakende die gebruik en verenigbaarheid met ander produkte.

GEBRUIKSAANWYSINGS

- Meng die verlangde hoeveelheid ComCat® MAX WP vooraf in 5-10 L water. Voeg die verlangde hoeveelheid Aqua-Wet® by teen 10ml/100L water (water volume van toedieningstenk) by ComCat® MAX WP mengsel en roer deeglik. Voeg AnnGro® by laasgenoemde teen verlangde hoeveelheid (240 ml/ha) en laat mengsel staan vir 30 minute voordat dit by toedieningstenk gevoeg word. Voeg ZumSil® teen verlangde hoeveelheid by tot toedieningstenk. Verseker deeglike mengaksie voor en gedurende toediening. Vir beste resultate moet nie dat mengsel oornag staan nie. Die gebruik van geskikte adjuvant word aanbeveel vir beter resultate.

TOEDIENINGSMETODE

Gewas	Tyd van toediening	ComCat® MAX WP dosis (g/ha)	ZumSil® dosis (L/ha)
Mielies	9-15 blare ontvou	50	1
Koring	Groeistadium 8 of 16 (Joubert)	100	1

WAARSKUWING

Alhoewel hierdie produk omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en die effek daarvan beïnvloed kan word deur faktore soos abnormale grond, klimaats- en bergingstoestande, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing, of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van die toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.