



Reg. No. M33 Act. No. 36 of 1947

ComCat® WP

500g

A BIOSTIMULANT

Optimizing the growth and yield of crops.

*N BIOSTIMULANT

Optimaliseer die groei en opbrengs van gewasse.

FERTILIZER GROUP

3

MISSTOF GROEP

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

2,4- Epibrassinolied _____ 0.3 mg/kg _____ 2,4-Epibrassinolide



 **agraforum**

REGISTRATION HOLDER / REGISTRASIE HOUER

AgraForum South Africa (Pty) Ltd | Reg. no: 2004/022326/07

P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330 | www.agraforum-sa.com



Batch Number | Date of manufacturing
Lot Nommer | Datum van vervaardiging



GENERAL

- ComCat® is water-soluble and non-toxic to plants when applied as directed.
- For best results apply ComCat® according to recommendations.
- Applied as a foliar spray.
- Before using ComCat®, consult an agriculturalist or the manufacturer on the use and compatibility with other products.

PRECAUTIONS

- It's misuse may cause damage to crops.
- Mild irritant to eyes. Wear protective clothing. In case of eye contact, wash with clean running water.
- Do not eat, drink or smoke whilst mixing or applying.
- Store in a cool (below 24°C) and dry place.

DIRECTIONS FOR USE

- Premix the prescribed amount of ComCat® in 0.5-4 L of water and mix thoroughly before application to spray tank. Ensure thorough mixing before and during application. For best results, do not let prepared mixture stand overnight. Apply to foliage in adequate spray volume to ensure good crop coverage. The use of the correct adjuvant is recommended for better uptake.

METHOD OF APPLICATION

Crop	Time of Application	Dosage (g/ha)
Maize	5-8 leaves unfolded	100
Wheat	Growth stage 5-11 (Joubert)	200
Groundnuts	Beginning of pod development	200

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

ALGEMEEN

- ComCat® is oplosbaar in water en nie-toksies vir plante wanneer toegedien soos voorgeskryf.
- Vir beste resultate dien ComCat® toe volgens aanbeveling.
- Toegedien as 'n blaarbespuiting.
- Voordat ComCat® gebruik word, raadpleeg 'n landboukundige of die vervaardiger rakende die gebruik en verenigbaarheid met ander produkte.

VOORSORGMATREËLS

- Wangebruik kan gewasse beskadig.
- Mag irriterend wees vir die oë. Dra beskermende klere. In die geval van oogkontak, was met skoon lopende water.
- Moet nie eet, drink of rook gedurende meng of aanwending nie.
- Berg in 'n koel (laer as 24°C) en droë plek.

GEBRUIKSAANWYSINGS

- Meng die verlangde hoeveelheid ComCat® vooraf in 0.5-4 L water en meng deeglik voor byvoeging tot spuitenk. Verseker deeglike mengaksie voor en gedurende toediening. Vir beste resultate, moet nie dat mengsel oornag staan nie. Dien toe aan blare teen genoegsame spuitvolume om goeie gewasbedekking te verseker. Die gebruik van geskikte adjuvant word aanbeveel vir beter opname.

TOEDIENINGSMETODE

Gewas	Tyd van toediening	Dosis (g/ha)
Mielies	5-8 blare ontvou	100
Koring	Groeistadium 5-11 (Joubert)	200
Grondbone	Begin van peulvorming	200

WAARSKUWING

Alhoewel hierdie produk omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie aangesien die werking en die effek daarvan beïnvloed kan word deur faktore soos abnormale grond, klimaats- en bergingstoestande, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing, of vir nadelige effek op mens of dier of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings na te kom of as gevolg van die ontstaan van die toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.