AgraVito R

Read the label before use Keep out of reach of children and animals

POWDER PLANT NUTRIENT A fertilizer containing macro nutrients

FERTILIZER GROUP 1





Signal word: DANGER

Hazard Statement:

- Causes severe skin burns and eve damage
- · May intensify fire: oxidiser

Precautionary Statement:

- Wash hands and face thoroughly after handling
- Wear protective gloves / clothing / eye protection / face protection
- · Keep away from heat, hot surface, sparks, open flames and other ignition sources
- No smoking

PRODUCT COMPOSITION

N: 129 a/ka P: 96 g/kg K: 201 a/ka S: 38 g/kg



AgraForUm South Africa (Pty) Ltd | Reg. no: 2004/022326/07 P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330



WARNING:

- Causes severe skin burns and eye damage.
- · May intensify fire: oxidiser.
- Keep away from heat, hot surface, sparks, open flames and other ignition sources. No smoking.
- · Do not store mixed product.
- Might be corrosive to metals, pH = 1.45.

PRECAUTIONS:

- Do not breathe dust/fume/gas/mist/vapors/spray.
- · Wash hands and face thoroughly after handling.
- Wear protective gloves/protective clothing/eye protection/face protection.
- . IF SWALLOWED: Rinse mouth, Do NOT induce vomiting.
- Take off immediately all contaminated clothing. Immediately rinse with water for several minutes.
- · Wash contaminated clothing before reuse.
- IF INHALED: Remove person to fresh air and keep comfortable for breathing.
- IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

RELEVANT SUBSTANCE:

Substance	Concentration
Urea	> 40 %
Potassium Nitrate	< 40 %

GENERAL

- AgraVito R™ is a fertilizer intended to be applied via dripper irrigation system to crops during the reproductive growth period.
- Do not apply to plants when plants are wilted or under stress.

APPLICATION RATES

Crops	Application rate	Remarks
Row crops	Via dripper irrigation system 10-100 kg/ha	Apply from 3 leaf stage and repeat application as required (at least 1-2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Vegetable crops	Via dripper irrigation system 10-100 kg/ha	Apply from 3 leaf stage and repeat application as required (at least 1-2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Fruit Tree crops	Via dripper irrigation system 10-100 kg/ha	Apply every 1-2 weeks during season as required based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.