



Read the label before use
Keep out of reach of children and animals

LIQUID PLANT NUTRIENT A fertilizer containing a micronutrient

FERTILIZER GROUP

2



Signal word: WARNING

Hazard Statements:

- Harmful if swallowed
- Causes skin irritation
- Causes serious eye irritation

Precautionary Statement:

- Wash face, hands and any exposed skin thoroughly after handling
- Do not eat, drink or smoke when using this product

PLANT NUTRIENT CONTENT

Fe: 72 g/kg
SG: 1.2 @ (20 °C)

NET VOLUME

10 L



REGISTRATION HOLDER

AgraForUm South Africa (Pty) Ltd, Reg. no: 2004/022326/07
P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330
mail@agraforum-sa.com

Batch number | Date of manufacturing



WARNING

- Harmful if swallowed
- Causes skin irritation
- Causes serious eye irritation

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTION

- Wash face, hands and any exposed skin thoroughly after handling.
- Do not eat, drink or smoke when using this product.
- Avoid release to the environment.
- Wear protective gloves/protective clothing/eye protection/face protection.
- IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.
- IF ON SKIN: Wash with plenty of soap and water.
- Specific treatment (see supplemental first aid instructions on this label).
- If skin irritation occurs: Get medical advice/attention.
- Take off contaminated clothing and wash before reuse.
- IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.
- Immediately call a POISON CENTER or doctor/physician.
- Rinse mouth.
- Dispose of contents/container according to local regulations.
- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.

GENERAL

- Ironman is a fertilizer intended as a supplement to normal fertilizer programs.
- Foliar spraying should preferably be done during the cooler periods of the day (ideal 20°C; < 40°C), as well as when the relative humidity is at its highest.
- Wind speed should be < 8 km/h.
- Preferably spray water pH 5-6.
- Do not apply to plants when plants are wilted or under stress.

COMPATIBILITY

- Always verify compatibility when mixed with other agricultural remedies (jar test as well as first spraying small area to check for phytotoxicity).
- Always test the product first on small area to check for phytotoxicity especially if higher rates are used.

RELEVANT SUBSTANCE

Ingredient		% Composition
Iron sulphate heptahydrate		< 40 %
Citric Acid		< 1.5 %

APPLICATION RATES

Crop	Application rate	Remarks
Row crops	Foliar spray 200-750 ml/100 L water. Via irrigation system 5-25 L/ha. (Minimum dilution rate 1L Ironman:250L water)	Use minimum 200 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 10-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop.
Vegetable crops	Foliar spray 200-750 ml/100 L water. Via irrigation system 5-25 L/ha. (Minimum dilution rate 1L Ironman:200L water)	Use minimum 300 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Repeat application as required (at least 10-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop.
Fruit Tree crops	Foliar spray 200-750 ml/100 L water. (maximum 10 L/ha) Via irrigation system 15-25 L/ha. (Minimum dilution rate 1L Ironman:250L water)	Apply as required based on soil and/or leaf analysis. Repeat application as required (at least 10-14 days apart). Do not exceed the nutrient requirement of the crop.