

Reg. No. B5564 Act No.36 of 1947

LiQuiDo Base

Read the label before use
Keep out of reach of children and animals

LIQUID PLANT NUTRIENT A fertilizer containing micronutrients

FERTILIZER GROUP **2**



Signal word: DANGER

Hazard Statements:

- May damage fertility or the unborn child
<Perinatal mortality>

Precautionary Statements:

- Obtain special instructions before use
- Do not handle until all safety precautions have been read and understood

ACTIVE INGREDIENTS

Mn: 22 g/kg
Zn: 16 g/kg
Cu: 6.3 g/kg
B: 6.7 g/kg
SG @ 20°C: 1.22

NET VOLUME

25 L



REGISTRATION HOLDER

AgraForUm South Africa (Pty) Ltd | Reg. no: 2004/022326/07
P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330
mail@agraforum-sa.com

Batch number | Date of manufacturing



WARNING

- May damage fertility or the unborn child <Perinatal mortality>

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTION

- Obtain special instructions before use.
- Do not handle until all safety precautions have been read and understood.
- Use personal protective equipment as required.
- IF exposed or concerned: Get medical advice/attention.
- Store locked up.
- Dispose of contents/container according to local regulations.
- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.

GENERAL

- **LiQuiDo BASE** is a fertilizer intended as a supplement to normal fertilizer programs.
- Foliar spraying should preferably be done during the cooler periods of the day (ideal 20°C; < 40°C), as well as when the relative humidity is at its highest.
- Wind speed should be < 8km/h.
- Do not apply to plants when plants are wilted or under stress.

RELEVANT SUBSTANCE

Ingredient	Composition %
Copper EDTA 150	1-10%
Boric acid 170	1-10%

COMPATIBILITY

- Always verify compatibility when mixed with other products (jar test as well as first spraying a small area to check for phytotoxicity).
- Always test the product first on a small area to check for phytotoxicity especially if higher rates are used.

APPLICATION RATES

Crop	Application rate	Remark
Soil application		
General	5 L-25 L/ha	Apply as required based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop.
Foliar application		
Row crops	Foliar spray 200 ml-1 L/100 L water. (maximum 5 L/ha)	Use minimum 200 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 500 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 7-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop.
	Via irrigation system 5-20 L/ha. (maximum 20 L/ha)	
Vegetable crops	Foliar spray 200 ml-1 L/100 L water. (maximum 5 L/ha)	Use minimum 300 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 500 L/ha). Repeat application as required (at least 7-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop.
	Via irrigation system 5-20 L/ha. (maximum 20 L/ha)	
Fruit Tree crops	Foliar spray 200 ml-1 L/100 L water. (maximum 10 L/ha)	Apply as required based on soil and/or leaf analysis. Repeat application as required (at least 7-14 days apart). Do not exceed the nutrient requirement of the crop.
	Via irrigation system 5-20 L/ha. (maximum 20 L/ha)	