



Reg. No. K10651 Act. No. 36 of 1947

HumiFol™

10L

A LIQUID PLANT NUTRIENT
A fertilizer containing nickel and selenium.

'N VLOEIBARE PLANTVOEDINGSTOF
'n Kunsmisproduk wat nikkel en selenium bevat.

FERTILIZER GROUP

2

MISSTOF GROEP

ACTIVE INGREDIENT / AKTIEWE BESTANDDEEL

Nickel _____	182 mg/kg	_____	Nikkel
Selenium _____	13 mg/kg	_____	Selenium
SG _____	1,24	_____	SG



REGISTRATION HOLDER / REGISTRASIE HOUER

AgraForum South Africa (Pty) Ltd | Reg. no: 2004/022326/07

P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330 | www.agraforum-sa.com

Batch Number | Date of manufacturing
Lot Nommer | Datum van vervaardiging



GENERAL

- HumiFol™ is a fertilizer intended as a supplement to normal fertilizer programs.
- Foliar spraying should preferable be done during the cooler periods of the day (ideal 20 °C; < 40°C), as well as when the relative humidity is on it's highest.
- Wind speed should be < 8 km/h.
- Do not apply to plants when plants are wilted or under stress.
- Please use a buffering agent to buffer pH between 5 and 6.
- Please use a suitable water-soluble wetting agent and penetrant (such as AnnGro™) to improve wetting, sticking and penetrating properties.
- Preferable use wide angle hollow cone spray nozzles.
- HumiFol™ is very hygroscopic.

PRECAUTION

- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.

COMPATIBILITY

- Always verify compatibility when mixed with other agricultural remedies (jar test as well as first spraying small area to check for phytotoxicity).
- Always test product first on small area to check for phytotoxicity especially if higher rates are used.
- Can be used in combination with ZincNiSeI™ and/or AgraActivator 5™.

APPLICATION RATES

Crop	Dosage rate	Remarks
Row crops	Foliar spray 50-500 ml/100 L	Use minimum 200 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.
Vegetable crops & Fruit trees	Foliar spray 50-500 ml/100 L water	Use minimum 300 L water/ha for foliar application. Better results might be obtained if a higher water rate is used. Repeat application as required (at least 2 weeks apart) based on soil and/or leaf analysis. Do not exceed nutrient requirement of the crop.

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.