



Reg. No. B5663 Act. No. 36 of 1947

Ironman™

10L

A LIQUID PLANT NUTRIENT
A fertilizer containing a micronutrient.

'N VLOEIBARE PLANTVOEDINGSTOF
'n Bemestingstof wat 'n mikro-elemente bevat.

FERTILIZER GROUP

2

MISSTOF GROEP

PLANT NUTRIENT CONTENT / PLANT VOEDINGSTOF INHOUD

Fe: 72 g/kg
SG 1.2 @ (20 °C)



 **agraforum**

REGISTRATION HOLDER / REGISTRASIE HOUER
AgraForUm South Africa (Pty) Ltd | Reg. no: 2004/022326/07
P.O. Box 38910, Langenhoven Park, Bloemfontein, 9330 | mail@agraforum-sa.com

Batch Number | Date of manufacturing
Lot Nommer | Datum van vervaardiging



PRECAUTION

- Always wear appropriate personal protective equipment.
- Keep out of reach of children.
- Do not store mixed product.

GENERAL

- Ironman is a fertilizer intended as a supplement to normal fertilizer programs.
- Foliar spraying should preferably be done during the cooler periods of the day (ideal 20 °C; < 40°C), as well as when the relative humidity is on its highest.
- Wind speed should be < 8 km/h.
- Preferable spray water pH 5-6.
- Do not apply to plants when plants are wilted or under stress.

COMPATIBILITY

- Always verify compatibility when mixed with other agricultural remedies (jar test as well as first spraying a small area to check for phytotoxicity).
- Always test the product first on a small area to check for phytotoxicity especially if higher rates are used.

APPLICATION RATES

| Crop | Application rate | Remarks |
|------------------|---|---|
| Row crops | Foliar spray 200-750 ml/100 L water. Via irrigation system 5-25 L/ha. (Minimum dilution rate 1L Ironman: 250L water) | Use minimum 200 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Apply at 3-8 leaf stage and repeat application as required (at least 10-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop. |
| Vegetable crops | Foliar spray 200-750 ml/100 L water. Via irrigation system 5-25 L/ha. (Minimum dilution rate 1L Ironman: 200L water) | Use minimum 300 L water/ha for foliar application. Better results might be obtained if a higher water rate is used (up to 400 L/ha). Repeat application as required (at least 10-14 days apart) based on soil and/or leaf analysis. Do not exceed the nutrient requirement of the crop. |
| Fruit tree crops | Foliar spray 200-750 ml/100 L water (maximum 10 L/ha) Via irrigation system 15-25 L/ha (Minimum dilution rate 1L Ironman: 250L water) | Apply as required based on soil and/or leaf analysis. Repeat application as required (at least 10-14 days apart). Do not exceed the nutrient requirement of the crop. |

WARNING

Although this product has been extensively tested under a large variety of conditions, the manufacturer does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, compatibility with other substances not indicated on the label as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal or for lack of performance of the product concerned due to failure of the user to follow label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.